

Violence During Pregnancy Among Women With or at Risk for HIV Infection

Linda J. Koenig, PhD, Daniel J. Whitaker, PhD, Rachel A. Royce, PhD, MPH, Tracey E. Wilson, PhD, Michelle R. Callahan, PhD, and M. Isabel Fernandez, PhD, for the Perinatal Guidelines Evaluation Project Group

Conservative estimates suggest that 4% to 8% of women in the United States experience violence during pregnancy.^{1,2} Several factors suggest that risk for violence may be even higher for pregnant women with HIV infection. Women with HIV are at increased risk for violence relative to the general population,^{3,4} perhaps because demographic and behavioral factors associated with HIV (e.g., poverty, drug use, bartering sex) also increase a woman's exposure to violence.^{5–9} In addition, some HIV-infected women may be at risk for violence when their positive serostatus is disclosed.^{10–15} Because a large proportion of HIV infections in women are diagnosed through routine prenatal screening,^{16–19} many disclosures may occur during pregnancy. Moreover, modifications of obstetric and postpartum care to prevent perinatal transmission (e.g., additional medications, formula feeding)^{20,21} may make it more difficult for women who are pregnant or have recently given birth to keep their serostatus private.

Data collected as part of the Perinatal Guidelines Evaluation Project—HIV and Pregnancy Study were used to examine the prevalence of violence during pregnancy among women with HIV and to ascertain if, and in what way, violence might be related to HIV serostatus. Specifically, seronegative women were matched to seropositive women on a number of demographic and HIV risk characteristics also associated with risk for violence. Owing to concerns that widespread prenatal screening for HIV might inadvertently increase women's risk for negative social consequences such as violence, the impact of receiving an HIV diagnosis during pregnancy was assessed. Finally, the association between recent violence and current partnership category was assessed and, for HIV-infected women, violence associated specifically with partner serostatus disclosure was examined.

Objectives. This study estimated the prevalence of violence during pregnancy in relation to HIV infection.

Methods. Violence, current partnerships, and HIV risk behaviors were assessed among 336 HIV-seropositive and 298 HIV-seronegative at-risk pregnant women.

Results. Overall, 8.9% of women experienced recent violence; 21.5% currently had abusive partners. Violence was experienced by women in all partnership categories (range = 3.8% with nonabusive partners to 53.6% with physically abusive partners). Neither experiencing violence nor having an abusive partner differed by serostatus. Receiving an HIV diagnosis prenatally did not increase risk. Disclosure-related violence occurred, but was rare.

Conclusions. Many HIV-infected pregnant women experience violence, but it is not typically attributable to their serostatus. Prenatal services should incorporate screening and counseling for all women at risk for violence. (*Am J Public Health.* 2002;92:367–370)

METHODS

Between October 1996 and October 1998, pregnant women receiving prenatal care (336 HIV-infected and 298 HIV-uninfected women) were recruited from health departments and clinics in Brooklyn, NY (n=224), Connecticut (n=55), Miami, Fla (n=220), and North Carolina (n=135), either directly or through their providers. All HIV-infected women at participating clinics were eligible. Eligible uninfected women (those receiving services at the clinic and testing negative for HIV during the current pregnancy) were screened and matched ($\pm 5\%$) to HIV-infected women within the same state according to the frequency of the following: sexual transmission risk behavior (ever used crack, had sex with a man known to use or suspected of using intravenous drugs, or traded sex for drugs or money), intravenous drug use, race/ethnicity, and timing of entry into prenatal care. (The cohort and design have been previously described.²²)

Demographic, behavioral, and HIV-related information was collected from women through interviews (in English, Spanish, or Haitian Creole) at 24 weeks or more into the pregnancy. Violence during pregnancy, referred to as recent violence, was assessed by

asking women whether they had been “beaten, physically attacked, or physically abused” or “sexually attacked, raped, or sexually abused” by anybody during the past 6 months. Relationship with an abusive partner was categorized by asking women with a main male partner how frequently, when upset, does he “physically abuse or hurt you” or “verbally or emotionally abuse you.” Women answering “always,” “often,” or “sometimes” were considered to have an abusive main male partner. Thus, we measured the proportion of women who experienced acts of violence during the specified period (regardless of perpetrator), or who currently had an abusive intimate partner (regardless of whether a violent act occurred during pregnancy). HIV-infected women were asked, “Does your partner know about your HIV infection?” and “Did anything bad happen when he found out about your HIV?”

RESULTS

Demographic and behavioral characteristics of the HIV-infected sample reflected those of the broader HIV epidemic among women and, with few exceptions, were similar to those of uninfected women (Table 1). Recent violence was reported by 8.9% of women. All

TABLE 1—Demographic, Behavioral, and Violence–Abuse Characteristics for All Women and by HIV Serostatus: Perinatal Guidelines Evaluation Project—HIV and Pregnancy Study

Variable	Total Sample (n = 634)	HIV+ (n = 336)	HIV- (n = 298)	P
Demographic/Behavioral Characteristics				
Mean age, y	27.8	28.3	27.2	.04
Race/ethnicity, ^a %				.50
White	7.3	5.9	8.7	
Hispanic	19.9	19.4	20.5	
Black (non-Hispanic)	69.7	71.1	68.1	
Other	3.2	3.6	2.3	
Education, %				.28
< High school	53.6	55.9	51.2	
High school	24.2	24.5	23.9	
> High school	22.2	19.6	24.9	
Monthly household income, %				.52
< \$1000	76.0	75.0	77.2	
≥ \$1000	24.0	25.0	22.8	
Employment outside the home, %				.74
Yes	21.5	21.0	22.0	
No	78.5	79.0	78.0	
Income sources, %				
Medicaid	86.6	86.0	87.2	.65
WIC	79.4	81.4	77.2	.19
Food stamps	49.8	57.8	40.9	.001
Disability	13.9	21.3	5.7	.001
Public assistance/welfare	29.7	33.2	25.8	.04
Family/partners	74.1	69.8	78.9	.01
No. of times moved in past year (mean)	1.1	0.97	1.24	.05
Gestational age at entry into prenatal care ^a				.61
≥ 20 wk	14.7	14.0	15.4	
< 20 wk	85.3	86.0	84.6	
Sex risk (crack, sex with intravenous drug using male, barter sex), ^a %				.41
Yes	39.3	40.8	37.6	
No	60.7	59.2	62.4	
Drug risk (intravenous drug use ever), ^a %				.09
Yes	4.9	6.3	3.4	
No	95.1	93.7	96.6	
Recent Violence and Partnership Category				
Experienced recent physical or sexual violence, ^b %				.13
Yes	8.9	7.3	10.7	
No	91.1	92.7	89.3	
Has a main male partner, ^c %				.003
Yes	81.8	77.6	86.6	
No	18.2	22.4	13.4	
Characterization of main male partner, ^d %				
Physically abusive	5.5	5.1	5.8	.72
Verbally/emotionally abusive	16.0	14.5	17.4	.36
Nonabusive	78.6	80.4	76.7	.31

Continued

but 1 woman who reported sexual violence also reported physical violence. Most women had a main male partner; 21.5% described their partner as physically or verbally/emotionally abusive. Violence was significantly associated with partnership category (highest for physically abusive, lowest for nonabusive, intermediate for verbally/emotionally abusive or no partner), but almost three fourths of the women who experienced violence were not currently in a relationship with a physically abusive partner (Table 2).

Neither violence nor having an abusive partner differed according to serostatus (Table 1). The proportion of women reporting violence was not higher among the 142 seropositive women diagnosed with HIV during the current pregnancy (5.8%) than among seronegative women (10.7%) or seropositive women with a prior HIV diagnosis (9.4%). Of the 260 HIV-infected women with main male partners, only 206 (79.2%) said that their partner knew their serostatus. Of these, 20 (9.7%) indicated that something bad happened when he found out. One woman was physically assaulted by her partner; 1 woman physically assaulted her partner. Other common negative outcomes included anger, depression, or shock (n=9) and relationship problems or terminations (n=6).

DISCUSSION

Overall, 8.9% of pregnant women in this study experienced recent violence. Although this is only slightly higher than average proportions previously reported,¹ it confirms that violence against pregnant women is a significant public health problem, endangering both the woman and her fetus.^{23–32} Extrapolating from the 6000 to 7000 yearly births to women in the United States with HIV,³³ we calculate that violence likely affects somewhere between 528 and 616 HIV-infected pregnant women each year. Although risk for violence did not differ by serostatus, we do not yet know if violence has a different impact on HIV-infected women, who are already at increased risk for adverse birth outcomes,^{22,34,35} and who must adhere to complex medication regimens during pregnancy.^{21,36}

Moreover, many more women may be at risk for violence and its health-related consequences, as 21.5% of the women in the study

TABLE 1—Continued

Experienced recent violence or had a physically or verbally/emotionally abusive partner, ^e %				.053
Yes	22.1	19.1	25.5	
No	77.9	80.9	74.5	

Note. WIC = Special Supplemental Nutrition Program for Women, Infants, and Children.

^aMatched in design.

^bRecent violence means the woman experienced violence in the prior 6 months. Six women for whom violence data were missing are excluded.

^cOne woman for whom male partner data were missing is excluded.

^dCategories are mutually exclusive and determined as a hierarchy. Five women with main male partners for whom data on partner abusiveness were missing are excluded.

^eOne woman for whom both violence and partner data were missing is excluded.

currently had an abusive primary partner. Not surprisingly, women with a physically abusive partner were most likely to experience recent violence. However, many women whose partner was not physically abusive, and even those without a partner, experienced violence. This reinforces the need for routine risk assessment in prenatal care settings and for increased awareness that women face risks from ex-partners^{5,37,38} and nonpartners,³⁹ as well as from current partners.

Consistent with other cohort studies of non-pregnant HIV-infected women in which the uninfected comparison women were drawn from demographically and behaviorally similar populations,^{3,4,40} neither violence nor having an abusive partner was associated with serostatus. Moreover, receiving a prenatal HIV diagnosis was not associated with elevated risk for violence. Taken together, these findings

suggest that violence against HIV-infected women is more likely related to the socioeconomic or behavioral contexts that characterize their lives than to serostatus itself. This is not to say that violence is never related to HIV.

One woman was physically assaulted when her partner learned her serostatus. Moreover, many HIV-infected women had not disclosed their serostatus to their partners, and we do not know to what extent this may have been due to fear of violence. Providers must be aware of these risks and concerns when addressing issues of disclosure.⁶ However, HIV-infected women's risk for violence may be best addressed by providing access to economic and social services (e.g., financial assistance, substance abuse treatment) that target conditions known to be associated with risk of violence. The prenatal care setting, with its multiple scheduled provider contacts, may

provide an important opportunity for identifying and referring women at risk.

These data represent a large and unique sample of HIV-infected and at-risk pregnant women. Nevertheless, certain limitations apply. Women in prenatal care may be at less risk for violence than those not in care. Moreover, by assessing only a 6-month period of time, some incidents of violence during pregnancy may have been missed. At the same time, because we enrolled women as early as 24 weeks into pregnancy, some incidents of violence may have occurred in the weeks just before conception. Future research will want to identify and address risk associated with specific pregnancy periods⁴¹ (e.g., preconception, postpartum) as well as the impact of partnership status on violence during those times. ■

About the Authors

Linda J. Koenig and Daniel J. Whitaker are with the Division of HIV/AIDS Prevention, National Center for HIV, STD, and TB Prevention, Centers for Disease Control and Prevention, Atlanta, Ga. At the time of the study, Rachel A. Royce was with the Department of Epidemiology, University of North Carolina at Chapel Hill. Tracey E. Wilson is with the Department of Preventive Medicine and Community Health, Health Science Center, State University of New York at Brooklyn. Michelle R. Callahan is with the Department of Epidemiology and Public Health, Yale School of Medicine, New Haven, Conn. At the time of this study, M. Isabel Fernandez was with the Department of Psychiatry and Behavioral Sciences, University of Miami School of Medicine, Miami, Fla.

Requests for reprints should be sent to Office of Communications, National Center for HIV, STD, and TB Prevention, Mail Stop E-06, Atlanta, GA 30333

This article was accepted April 4, 2001.

TABLE 2—Number and Proportion of Women Who Experienced Violence in Prior 6 Months, by Characterization of Current Male Partner

	Experienced Violence in Prior 6 mo						P (HIV+ /HIV- Comparison)
	Total Sample (n = 623)		HIV-Seropositive Women (n = 325)		HIV-Seronegative Women (n = 298)		
	No, n (%)	Yes, n (%)	No, n (%)	Yes, n (%)	No, n (%)	Yes, n (%)	
Characterization of current main male partner							
Physically abusive	13 (46.4)	15 (53.6) ^a	7 (53.8)	6 (46.2) ^a	6 (40.0)	9 (60.0) ^a	.46
Verbally/emotionally abusive	71 (86.6)	11 (13.4) ^b	33 (89.2)	4 (10.8) ^{b,c}	38 (84.4)	7 (15.6) ^b	.53
Nonabusive	383 (96.2)	15 (3.8) ^c	195 (97.5)	5 (2.5) ^c	188 (95.0)	10 (5.0) ^b	.18
No main male partner	100 (88.0)	15 (12.0) ^b	66 (88.0)	9 (12.0) ^b	34 (85.0)	6 (15.0) ^b	.65
Total	567 (91.0)	56 (9.0)	301 (92.6)	24 (7.4)	266 (89.3)	32 (10.7)	
Overall χ^2_3 for sample	85.6 (P<.0001)		38.5 (P<.0001)		11.1 (P< .001)		

^{a,b,c}Among the 6 pairwise comparisons within each sample (i.e., physically abusive vs verbally/emotionally abusive, physically abusive vs nonabusive, physically abusive vs no main male partner, verbally/emotionally abusive vs nonabusive, verbally/emotionally abusive vs no main male partner, and nonabusive vs no main male partner), proportions in a column not sharing the same superscript letter are significantly different at P < .008 (Bonferroni correction for 6 comparisons).

Contributors

L.J. Koenig conceived of the study, with collaboration from all authors, and drafted and finalized the manuscript. D.J. Whitaker analyzed the data with assistance from L.J. Koenig and R.A. Royce. D.J. Whitaker, R.A. Royce, T.E. Wilson, M.R. Callahan, and M.I. Fernandez contributed to the interpretation of the results and to revisions of the paper. R.A. Royce, T.E. Wilson, and M.I. Fernandez directed study activities in their respective cities.

Acknowledgments

The Perinatal Guidelines Evaluation Project was funded by the Centers for Disease Control and Prevention (CDC) through cooperative agreements U64CCU412273 (Duke), U64CCU112274 (Yale), U64CCU412294 (Miami), and U64CCU212267 (SUNY). All procedures and consent forms were approved by the Institutional Review Boards at the Centers for Disease Control and Prevention and each participating university and hospital. We are grateful to the members of the study staff who assisted with data collection and project management and to the women who participated in the study.

Members of the Perinatal Guidelines Evaluation Project Group include Howard Minkoff, MD, Jack DeHovitz, MD, and Tracey Wilson, PhD (New York); Rachel Royce, PhD, and Emmanuel Walter, MD (North Carolina); Mary Jo O'Sullivan, MD, and M. Isabel Fernandez, PhD (Florida); Jeannette Ickovics, PhD, and Kathleen Ethier, PhD (Connecticut); and Linda Koenig, PhD, R.J. Simonds, MD, Jan Moore, PhD, Margaret Lampe, RN, Daniel Whitaker, PhD, Daphne Cobb, MS, and Ken Dominguez, MD (CDC).

References

- Gazmararian JA, Lazorick S, Spitz AM, Ballard TJ, Saltzman LE, Marks JS. Prevalence of violence against pregnant women. *JAMA*. 1996;275:1915–1920.
- Tjaden P, Thoennes N. *Prevalence, Incidence, and Consequences of Violence Against Women: Findings From the National Violence Against Women Survey*. Washington, DC: US Dept of Justice, National Institute of Justice; 1998. US Dept of Justice publication NCJ 172837.
- Cohen M, Deamant C, Barkan S, et al. Domestic violence and childhood abuse in HIV-infected women and women at risk for HIV. *Am J Public Health*. 2000; 90:560–565.
- Vlahov D, Wientge D, Moore J, et al. Violence against women with or at risk for HIV infection. *AIDS Behav*. 1998;2:53–60.
- Bachman R, Saltzman LE. *Violence Against Women: Estimates From the Redesigned Survey*. Washington, DC: US Dept of Justice; 1995. US Dept of Justice publication NCJ 154348.
- Koenig LJ, Moore J. Women, violence and HIV: a critical evaluation with implications for HIV services. *Matern Child Health J*. 2000;4:103–109.
- Maman D, Campbell J, Sweat MD, Geilen AC. The intersections of HIV and violence: directions for future research and interventions. *Soc Sci Med*. 2000;50: 459–478.
- Zierler S. Hitting hard: HIV and violence against women. In: Goldstein N, Manlow J, eds. *The Gender Politics of HIV*. New York, NY: New York University Press; 1997:207–221.
- Zierler S. Reframing women's risk: social inequalities and HIV infection. *Annu Rev Public Health*. 1997; 18:401–436.
- Gielen AC, O'Campo P, Faden RR, Eke A. Women's disclosure of HIV status: experiences of mistreatment and violence in an urban setting. *Women Health*. 1997;25:19–31.
- Gielen AC, McDonnell K, Burke J, O'Campo P. Women's lives after an HIV positive diagnosis: disclosure and violence. *Matern Child Health J*. 2000;4: 111–120.
- North RL, Rothenberg KH. Partner notification and the threat of domestic violence against women with HIV infection. *N Engl J Med*. 1993;329:1194–1196.
- Rothenberg KH, Paskey SJ, Reuland MM, Zimmerman SI, North RL. Domestic violence and partner notification: implications for treatment and counseling of women with HIV. *J Am Med Womens Assoc*. 1995;50: 87–93.
- Rothenberg KH, Paskey SJ. The risk of domestic violence and women with HIV infection: implications for partner notification, public policy, and the law. *Am J Public Health*. 1995;85:1569–1576.
- Zierler S, Cunningham WE, Andersen R, et al. Violence victimization after HIV infection in a US probability sample of adult patients in primary care. *Am J Public Health*. 2000;90:208–215.
- Centers for Disease Control and Prevention. Success in implementing public health service guidelines to reduce perinatal transmission of HIV—Louisiana, Michigan, New Jersey, and South Carolina, 1993, 1995, and 1996. *MMWR Morb Mortal Wkly Rep*. 1998;47:688–691.
- Centers for Disease Control and Prevention. Prenatal discussion of HIV testing and maternal HIV testing—14 states, 1996–1997. *MMWR Morb Mortal Wkly Rep*. 1999;48:401–404.
- Fernandez IM, Wilson TE, Ethier KA, Walter EB, Gay CL, Moore J. Acceptance of HIV testing during prenatal care. Perinatal Guidelines Evaluation Project. *Public Health Rep*. 2000;115:460–468.
- Royce R, Walter EB, Wilson T, Fernandez IM, Ickovics J, Simonds RJ. Barriers to universal prenatal HIV testing in 4 US sites in 1997. *Am J Public Health*. 2001;91:727–733.
- Centers for Disease Control and Prevention. Public Health Service recommendations for human immunodeficiency virus counseling and voluntary testing for pregnant women. *MMWR Morb Mortal Wkly Rep*. 1995;44(RR-7):1–15.
- Centers for Disease Control and Prevention. Public Health Service task force recommendations for the use of antiretroviral drugs in pregnant women infected with HIV-1 for maternal health and for reducing perinatal HIV-1 transmission in the United States. *MMWR Morb Mortal Wkly Rep*. 1998;47(RR-2):1–30.
- Ickovics JR, Ethier KA, Koenig LJ, Wilson TE, Walter EB, Fernandez MI, for the Perinatal Guidelines Evaluation Project. Infant birth weight among women with or at high risk for HIV infection: the impact of clinical, behavioral, psychosocial and demographic factors. *Health Psychol*. 2000;19:515–523.
- Newberger EH, Barkan SE, Lieberman ES, et al. Abuse of pregnant women and adverse birth outcome: current knowledge and implications for practice. *JAMA*. 1992;267:2370–2372.
- Connolly A, Katz VL, Bash KL, McMahon MJ, Hansen WF. Trauma and pregnancy. *Am J Perinatol*. 1997;14:331–336.
- Pak LL, Reece A, Chan L. Is adverse pregnancy outcome predictable after blunt abdominal trauma? *Am J Obstet Gynecol*. 1998;179:1140–1144.
- Petersen R, Gazmararian JA, Spitz AM, et al. Violence and adverse pregnancy outcomes: a review of the literature and directions for future research. *Am J Prev Med*. 1997;13:366–373.
- Poole GV, Martin JN, Perry KG, Griswold JA, Lambert CJ, Rhodes RS. Trauma in pregnancy: the role of interpersonal violence. *Am J Obstet Gynecol*. 1996;174: 1873–1876.
- Sandman CA, Wadhwa PD, Chicz-DeMet A, Dunkel-Schetter C, Porto M. Maternal stress, HPA activity, and fetal/infant outcome. *Ann NY Acad Sci*. 1997;814:266–275.
- Cooper RN, Goldenberg RL, Das A, et al. The preterm prediction study: maternal stress is associated with spontaneous preterm birth at less than thirty-five weeks gestation. National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. *Am J Obstet Gynecol*. 1996;175:1286–1292.
- Curry MA, Harvey SM. Stress related to domestic violence during pregnancy and infant birth weight. In: Campbell J, ed. *Empowering Survivors of Abuse: Health Care for Battered Women and Their Children*. Thousand Oaks, Calif: Sage Publications; 1998:98–108.
- McFarlane J, Parker B, Soeken K. Physical abuse, smoking, and substance use during pregnancy: prevalence, interrelationships, and effects on birth weight. *J Obstet Gynecol Neonatal Nurs*. 1996;25:313–320.
- Campbell J, Torres S, Ryan J, et al. Physical and nonphysical partner abuse and other risk factors for low birth weight among full term and preterm babies. *Am J Epidemiol*. 1999;150:7114–7126.
- Lindgren ML, Steinberg S, Byers RH. Epidemiology of HIV/AIDS in children. *Pediatr Clin N Am*. 2000;47:1–20.
- Temmerman M, Plummer FA, Mirza NB, et al. Infection with HIV as a risk factor for adverse obstetrical outcome. *AIDS*. 1990;4:1087–1093.
- Johnstone FD, Raab GM, Hamilton BA. The effect of human immunodeficiency virus infection and drug use on birth characteristics. *Obstet Gynecol*. 1996;88: 321–326.
- Wilson TE, Ickovics JR, Fernandez MI, Koenig LJ, Walter E. Self-reported zidovudine adherence among pregnant women living with human immunodeficiency virus infection in four US states. *Am J Obstet Gynecol*. 2001;184:1235–1240.
- Wilson MI, Daly M. Spousal homicide risk and estrangement. *Violence Vict*. 1993;8:3–16.
- Ellis D, DeKeseredy W. Rethinking estrangement, interventions, and intimate femicide. *Violence Against Women*. 1997;3:590–609.
- Grisso JA, Schwarz DF, Hirschinger N, et al. Violent injuries among women in an urban area. *N Engl J Med*. 1999;341:1899–1905.
- Zierler S, Witbeck B, Mayer K. Sexual violence against women living with or at risk for HIV infection. *Am J Prev Med*. 1996;12:304–310.
- Ballard TJ, Saltzman LE, Gazmararian JA, Spitz AM, Lazorick S, Marks JS. Violence during pregnancy: measurement issues. *Am J Public Health*. 1998;88: 274–276.